

$|V_{ub}|$ and $B \rightarrow \eta^{(\prime)}$ Form Factors in Covariant Light-Front Approach

Chuan-Hung Chen^{1,2*}, Yue-Long Shen^{3†} and Wei Wang^{4‡}

¹*Department of Physics, National Cheng-Kung University, Tainan 701, Taiwan*

²*National Center for Theoretical Sciences, Taiwan*

³*Institute of Physics, Academia Sinica, 115 Taipei, Taiwan*

⁴*Istituto Nazionale di Fisica Nucleare,
Sezione di Bari, Bari 70126, Italy*

(Dated: February 22, 2010)

Abstract

$B \rightarrow (\pi, \eta, \eta')$ transition form factors are investigated in the covariant light-front approach. With theoretical uncertainties, we find that $B \rightarrow (\pi, \eta, \eta')$ form factors at $q^2 = 0$ are $f_+^{(\pi, \eta, \eta')}(0) = (0.245_{-0.001}^{+0.000} \pm 0.011, 0.220 \pm 0.009 \pm 0.009, 0.180 \pm 0.008_{-0.007}^{+0.008})$ for vector current and $f_T^{(\pi, \eta, \eta')}(0) = (0.239_{-0.003-0.018}^{+0.002+0.020}, 0.211 \pm 0.009_{-0.015}^{+0.017}, 0.173 \pm 0.007_{-0.013}^{+0.014})$ for tensor current, respectively. With the obtained q^2 -dependent $f_+^\pi(q^2)$ and observed branching ratio (BR) for $\bar{B}_d \rightarrow \pi^+ \ell \bar{\nu}_\ell$, the V_{ub} is found as $|V_{ub}|_{LF} = (3.99 \pm 0.13) \times 10^{-3}$. As a result, the predicted BRs for $\bar{B} \rightarrow (\eta, \eta') \ell \bar{\nu}_\ell$ decays with $\ell = e, \mu$ are given by $(0.49_{-0.04-0.07}^{+0.02+0.10}, 0.24_{-0.02-0.03}^{+0.01+0.04}) \times 10^{-4}$, while the BRs for $D^- \rightarrow (\eta, \eta') \ell \bar{\nu}_\ell$ are $(11.1_{-0.6-0.9}^{+0.5+0.9}, 1.79_{-0.08-0.12}^{+0.07+0.12}) \times 10^{-4}$. In addition, we also study the integrated lepton angular asymmetries for $\bar{B} \rightarrow (\pi, \eta, \eta') \tau \bar{\nu}_\tau$: $(0.277_{-0.001-0.007}^{+0.001+0.005}, 0.290_{-0.000-0.003}^{+0.002+0.003}, 0.312_{-0.000-0.006}^{+0.004+0.005})$.

* Email: physchen@mail.ncku.edu.tw

† Email: shenyl@phys.sinica.edu.tw

‡ Email: wei.wang@ba.infn.it

One of the puzzles in exclusive B decays is the η' related processes. For example $B \rightarrow \eta' K$ was first observed by CLEO collaboration with the branching ratio (BR) $(89_{-16}^{+18} \pm 9) \times 10^{-6}$ [1] which is much larger than 30×10^{-6} estimated by the factorization ansatz. With more data accumulated, experimental uncertainties are pinning down and the world averages on the BR for $\eta^{(\prime)}$ production in B decays now are known as $\mathcal{B}(B^+ \rightarrow [\eta, \eta'] K^+) = [2.36 \pm 0.27, 71.1 \pm 2.6] \times 10^{-6}$, $\mathcal{B}(B^0 \rightarrow [\eta, \eta'] K^0) = [1.12_{-0.28}^{+0.30}, 66.1 \pm 3.1] \times 10^{-6}$ [2]. Clearly, the large BR for $B \rightarrow \eta' K$ is not smeared by statistic. To unravel the mystery, many novel solutions have been proposed, such as the intrinsic charm in η' [3], the gluonium state [4], the spectator hard scattering mechanism [5], the flavor-singlet component of η' [6] and enhanced chiral symmetry breaking effects [7]. Although it is believed that some exotic effects should be associated with $\eta^{(\prime)}$, it is difficult to specify where the novel effects should reside, since two-body hadronic B decays suffer from large uncertainties such as final state interactions.

Compared with nonleptonic B decays, semileptonic $\bar{B} \rightarrow \eta^{(\prime)} \ell \bar{\nu}_\ell$ and $\bar{B}_s \rightarrow \eta^{(\prime)} \ell^+ \ell^-$ decays are much cleaner and thus might be more helpful to explore the differences among various mechanisms. In particular, a sizable flavor-singlet component of $\eta^{(\prime)}$ predicts larger BRs for $\bar{B} \rightarrow \eta' \ell \bar{\nu}_\ell$ than the η modes, while the chiral symmetry breaking enhancement could give the reverse results [7]. Nevertheless, before one considers various possible novel effects on $\eta^{(\prime)}$, it is necessary to understand the BRs for $\bar{B} \rightarrow \eta^{(\prime)} \ell \bar{\nu}_\ell$ decays without these exotic effects. In our previous work [7], we used the perturbative QCD approach [8] to calculate the $B \rightarrow \eta^{(\prime)}$ form factors at large recoil; then the same whole spectrum as a function of invariant mass of $\ell \bar{\nu}_\ell$ for the form factors is assumed with that in the light-cone sum rules (LCSRs). Despite the predicted results for various branching ratios are consistent with the experimental data, it is meaningful to examine the same processes in other parallel frameworks. This is helpful to reduce the dependence on the treatments of the dynamics in transition form factors. The motif of this work is to employ another method to deal with the form factors: the covariant light-front (LF) approach [9, 10]. Since the predictions of $B \rightarrow \pi$ form factors in LF model match very well with those applied to the nonleptonic charmless B decays, it is worthy to understand what we can get the $B \rightarrow \eta^{(\prime)}$ form factors by this approach.

At the quark level, the $\bar{B} \rightarrow \eta^{(\prime)} \ell \bar{\nu}_\ell$ is induced by $b \rightarrow u \ell \bar{\nu}$ transition which will inevitably involve the $\bar{u}u$ component of the $\eta^{(\prime)}$ meson. Then the convenient mechanism for the $\eta - \eta'$




FIG. 1: Feynman diagram for the transition form factors, where the cross symbol in the diagram denotes the transition vertex.

mixing would be the quark flavor mixing scheme, defined by [11, 12]

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \eta_q \\ \eta_s \end{pmatrix}, \quad (1)$$

where $\eta_q = (u\bar{u} + d\bar{d})/\sqrt{2}$, $\eta_s = s\bar{s}$ and angle ϕ is the mixing angle. By the definition of $\langle 0 | \bar{q}' \gamma_\mu \gamma_5 q' | \eta_{q'}(p) \rangle = i f_{q'} p_\mu$ ($q' = q, s$), the masses of $\eta_{q,s}$ can be expressed by

$$m_{qq}^2 = \frac{\sqrt{2}}{f_q} \langle 0 | m_u \bar{u} i \gamma_5 u + m_d \bar{d} i \gamma_5 d | \eta_q \rangle, \quad m_{ss}^2 = \frac{2}{f_s} \langle 0 | m_s \bar{s} i \gamma_5 s | \eta_s \rangle. \quad (2)$$

Here, m_{qq} and m_{ss} are unknown parameters and their values can be obtained by fitting with the data. In terms of the quark-flavor basis, we see clearly that m_{qq} and m_{ss} are zero in the chiral limit. The advantage of the quark-flavor mixing scheme is: at the leading order in α_s only the quark transition from the B meson into the η_q component is necessary; while the other transitions like $B \rightarrow \eta_s$ are suppressed by α_s . The gluonic form factors (or referred to as flavor-singlet form factors) will be remarked later.

For calculating the transition form factors, we parameterize the hadronic effects as

$$\begin{aligned} \langle P(P'') | \bar{q}' \gamma^\mu b | \bar{B}(P') \rangle &= f_+^P(q^2) \left(P^\mu - \frac{P \cdot q}{q^2} q^\mu \right) + f_0^P(q^2) \frac{P \cdot q}{q^2} q_\mu, \\ \langle P(P'') | \bar{q}' i \sigma_{\mu\nu} q^\nu b | \bar{B}(P') \rangle &= \frac{f_T^P(q^2)}{m_B + m_P} [P \cdot q q_\mu - q^2 P_\mu], \\ \langle P(P'') | \bar{q}' \sigma_{\mu\nu} \gamma_5 b | \bar{B}(P') \rangle &= \frac{f_T^P(q^2)}{m_B + m_P} \epsilon_{\mu\nu\alpha\beta} P^\alpha q^\beta \end{aligned} \quad (3)$$

with $P_\mu = (P' + P'')_\mu$ and $q_\mu = (P' - P'')_\mu$. Since the light quarks in B -meson are u - and d -quark, the meson P could stand for π and η_q states.

In the covariant LF quark model, the transition form factors for $B \rightarrow P$ could be obtained by computing the lowest-order Feynman diagram depicted in Fig.1. Below we will adopt the

same notation as Ref. [9] and light-cone coordinate system for involved momenta, in which the components of meson momentum are read by $P' = (P'^-, P'^+, P'_\perp)$ with $P'^\pm = P'^0 \pm P'^3$. The relationship between meson momentum and the momenta of its constituent quarks is given by $P' = p'_1 + p_2$ and $P'' = p''_1 + p_2$ with p_2 being the spectator quark of initial and final mesons. Additionally, one can also express the quark momenta in terms of the internal variables $(x_i, p'_\perp)$ as

$$\tilde{p}_{1,2}^+ = x_{1,2} \tilde{P}^+, \quad \tilde{p}_{1,2\perp} = x_{1,2} \tilde{P}_\perp \pm \tilde{p}_\perp \quad (4)$$

with $x_1 + x_2 = 1$. Here, the notation with tilde could represent all momenta in the initial and final mesons.

In order to formulate the results of Fig. 1, the quark-meson-antiquark vertex for incoming and outgoing mesons are respectively chosen to be

$$\begin{aligned} i\Gamma'_P &= H'_P \gamma_5, \\ i(\gamma_0 \Gamma_P^\dagger \gamma_0) &, \end{aligned} \quad (5)$$

where H'_P is the covariant light-front wave function of the meson. Consequently, the amplitude for the loop diagram is straightforwardly written by

$$\mathcal{B}_{\mu(\mu\nu)}^{PP} = -i^3 \frac{N_c}{(2\pi)^4} \int d^4 p'_1 \frac{H'_P(H''_P)}{N'_1 N''_1 N_2} S_{\mu(\mu\nu)}^{PP}, \quad (6)$$

where $N_c = 3$ is the number of colors, $N_1^{(\prime\prime)} = p_1^{\prime(\prime)2} - m_1^{\prime(\prime)2}$, $N_2 = p_2^2 - m_2^2$.

$$\begin{aligned} S_\mu^{PP} &= 2p'_{1\mu} [M'^2 + M''^2 - q^2 - 2N_2 - (m'_1 - m_2)^2 - (m''_1 - m_2)^2 + (m'_1 - m''_1)^2] \\ &\quad + q_\mu [q^2 - 2M'^2 + N'_1 - N''_1 + 2N_2 + 2(m'_1 - m_2)^2 - (m'_1 - m''_1)^2] \\ &\quad + P_\mu [q^2 - N'_1 - N''_1 - (m'_1 - m''_1)^2], \\ S_{\mu\nu}^{PP} &= -\epsilon_{\mu\nu\alpha\beta} (-4m_2 p'_1{}^\alpha q^\beta + 2m''_1 p'_1{}^\alpha q^\beta - 2m''_1 P^\alpha p'_1{}^\beta \\ &\quad + 2m'_1 P^\alpha p'_1{}^\beta - 2m'_1 P^\alpha q^\beta + 2m'_1 p'_1{}^\alpha q_\beta), \end{aligned} \quad (7)$$

with the $M'(M'')$ being the mass of the incoming (outgoing) meson. As usual, the loop integral could be performed by the contour method. Therefore, except some separate poles appearing in the denominator, if the covariant vertex functions are not singular, the integrand is analytic. Thus, when performing the integration, the transition amplitude will pick

up the singularities from the anti-quark propagator so that the various pieces of integrand are led to be

$$\begin{aligned}
N_1'^{(n)} &\rightarrow \hat{N}_1'^{(n)} = x_1(M'^{(n)2} - M_0'^{(n)2}), \\
H_P'^{(n)} &\rightarrow h_P'^{(n)}, \\
S &\rightarrow \hat{S}, \\
\int \frac{d^4 p_1'}{N_1' N_1'' N_2} H_P' H_P'' S &\rightarrow -i\pi \int \frac{dx_2 d^2 p_1'}{x_2 \hat{N}_1' \hat{N}_1''} h_P' h_P'' \hat{S}.
\end{aligned} \tag{8}$$

We work in the $q^+ = 0$ frame and the transverse momentum of the quark in the final meson is given as $p_\perp'' = p_\perp' - x_2 q_\perp$. The new function of h_M' for initial meson is given by

$$h_P' = (M'^2 - M_0'^2) \sqrt{\frac{x_1 x_2}{N_c}} \frac{1}{\sqrt{2\widetilde{M}_0'}} \varphi_P', \tag{9}$$

with

$$\begin{aligned}
M_0'^2 &= (e_1' + e_2')^2 = \frac{p_\perp'^2 + m_1'^2}{x_1} + \frac{p_\perp'^2 + m_2'^2}{x_2}, \quad \widetilde{M}_0' = \sqrt{M_0'^2 - (m_1' - m_2')^2}, \\
e_i^{(\prime)} &= \sqrt{m_i^{(\prime)2} + p_\perp'^2 + p_z'^2}, \quad p_z' = \frac{x_2 M_0'}{2} - \frac{m_2'^2 + p_\perp'^2}{2x_2 M_0'},
\end{aligned} \tag{10}$$

where e_i can be interpreted as the energy of the quark or the antiquark, M_0' can be regarded as the kinetic invariant mass of the meson system and φ_P' is the LF momentum distribution amplitude for s -wave pseudoscalar mesons. The similar quantities associated with the outgoing meson can be defined by the same way.

After the contour integration, the valance antiquark is turned to be on mass-shell and the conventional LF model is recovered. The formulas of the form factors in the LF quark model shown in Eq. (7) would contain not only the terms proportional to P_μ and q_μ , but also the terms proportional to a null vector $\tilde{\omega} = (2, 0, \mathbf{0}_\perp)$. This vector is spurious, because it does not appear in the standard definition of Eq.(3), and spoils the covariance. In the literature, it is argued that this spurious factor can be eliminated by including the so-called zero-mode contribution, and a proper way to resolve this problem has been proposed in Ref. [9]. In this method, one should obey a series of special rules when performing the p^- integration. A manifest covariant result can be given with this approach, which is physically reasonable. Using Eqs. (7)–(9) and taking the advantage of the rules in Ref. [9, 10], the $B \rightarrow P$ form

factors are straightforwardly obtained by

$$\begin{aligned}
f_+^P(q^2) &= \frac{N_c}{16\pi^3} \int dx_2 d^2 p'_\perp \frac{h'_P h''_P}{x_2 \hat{N}'_1 \hat{N}''_1} \left[x_1 (M_0'^2 + M_0''^2) + x_2 q^2 \right. \\
&\quad \left. - x_2 (m'_1 - m''_1)^2 - x_1 (m'_1 - m_2)^2 - x_1 (m''_1 - m_2)^2 \right], \\
f_-^P(q^2) &= \frac{N_c}{16\pi^3} \int dx_2 d^2 p'_\perp \frac{2h'_P h''_P}{x_2 \hat{N}'_1 \hat{N}''_1} \left\{ -x_1 x_2 M'^2 - p_\perp'^2 - m'_1 m_2 + (m''_1 - m_2)(x_2 m'_1 + x_1 m_2) \right. \\
&\quad \left. + 2 \frac{q \cdot P}{q^2} \left(p_\perp'^2 + 2 \frac{(p'_\perp \cdot q_\perp)^2}{q^2} \right) + 2 \frac{(p'_\perp \cdot q_\perp)^2}{q^2} - \frac{p'_\perp \cdot q_\perp}{q^2} [M''^2 - x_2 (q^2 + q \cdot P) \right. \\
&\quad \left. - (x_2 - x_1) M'^2 + 2x_1 M_0'^2 - 2(m'_1 - m_2)(m'_1 + m''_1)] \right\}, \\
f_T^P(q^2) &= (M' + M'') \frac{N_c}{16\pi^3} \int dx_2 d^2 p'_\perp \frac{h'_P h''_P}{x_2 \hat{N}'_1 \hat{N}''_1} \left[x_1 (2m_2 - m'_1 - m''_1) + 2m'_1 \right. \\
&\quad \left. - 2(m'_1 - m''_1) \left(\frac{x_1}{2} - \frac{p'_\perp \cdot q_\perp}{q^2} \right) \right], \tag{11}
\end{aligned}$$

where the relation of $f_-^P(q^2)$ to $f_0^P(q^2)$ can be read by

$$f_0^P(q^2) = f_+^P(q^2) + \frac{q^2}{m_B^2 - m_P^2} f_-^P(q^2). \tag{12}$$

Clearly, one has $f_+^P(0) = f_0^P(0)$.

After we obtain the formulae for the $B \rightarrow P$ transition form factors, the direct application is the exclusive semileptonic $\bar{B} \rightarrow P \ell \bar{\nu}_\ell$ decays. The effective Hamiltonian for $b \rightarrow u \ell \bar{\nu}_\ell$ in the standard model (SM) is given by

$$H_{\text{eff}} = \frac{G_F V_{ub}}{\sqrt{2}} \bar{u} \gamma_\mu (1 - \gamma_5) b \bar{\ell} \gamma^\mu (1 - \gamma_5) \nu_\ell. \tag{13}$$

Although these decays are tree processes, however, if we can understand well the form factors, there still have the chance to probe the new physics in these semileptonic decays [14, 15]. Hence, the decay amplitude for $\bar{B} \rightarrow P \ell \bar{\nu}_\ell$ is written as

$$M(\bar{B} \rightarrow P \ell \bar{\nu}_\ell) = \langle \ell \bar{\nu}_\ell P | H_{\text{eff}} | \bar{B} \rangle = \frac{G_F V_{ub}}{\sqrt{2}} \langle P | \bar{u} \gamma_\mu (1 - \gamma_5) b | \bar{B} \rangle \bar{\ell} \gamma^\mu (1 - \gamma_5) \nu_\ell. \tag{14}$$

To calculate the differential decay rates, we choose the coordinates of various particles as follows

$$\begin{aligned}
q^2 &= (\sqrt{q^2}, 0, 0, 0), \quad p_B = (E_B, 0, 0, |\mathbf{p}_P|), \\
p_P &= (E_P, 0, 0, |\mathbf{p}_P|), \quad p_\ell = (E_\ell, |\mathbf{p}_\ell| \sin \theta, 0, |\mathbf{p}_\ell| \cos \theta), \tag{15}
\end{aligned}$$

where $E_P = (m_B^2 - q^2 - m_P^2)/(2\sqrt{q^2})$, $|\mathbf{p}_P| = \sqrt{E_P^2 - m_P^2}$, $E_\ell = (q^2 + m_\ell^2)/(2\sqrt{q^2})$ and $|\mathbf{p}_\ell| = (q^2 - m_\ell^2)/(2\sqrt{q^2})$. It is clear that θ is defined as the polar angle of the lepton momentum relative to the moving direction of the B -meson in the q^2 rest frame. With Eqs. (14) and (15), the differential decay rate for $\bar{B} \rightarrow P\ell\bar{\nu}_\ell$ as a function of q^2 and θ can be derived by

$$\begin{aligned} \frac{d\Gamma_P}{dq^2 d\cos\theta} &= \frac{G_F^2 |V_{ub}|^2 m_B^3}{2^8 \pi^3} \sqrt{(1-s+\hat{m}_P^2)^2 - 4\hat{m}_P^2} \left(1 - \frac{\hat{m}_\ell^2}{s}\right)^2 \\ &\quad \times [\Gamma_1^P + \Gamma_2^P \cos\theta + \Gamma_3^P \cos^2\theta], \\ \Gamma_1^P &= \hat{P}_P^2 f_+^{P2}(q^2) + (1 - \hat{m}_P^2)^2 \frac{\hat{m}_\ell^2}{s} f_0^{P2}(q^2), \\ \Gamma_2^P &= 2 \frac{\hat{m}_\ell^2}{s} \hat{P}_P (1 - \hat{m}_P^2) f_+^P(q^2) f_0^P(q^2), \\ \Gamma_3^P &= -\hat{P}_P^2 f_+^{P2}(q^2) + \frac{\hat{m}_\ell^2}{s} \hat{P}_P^2 f_+^{P2}(q^2), \end{aligned} \quad (16)$$

where $s = q^2/m_B^2$, $\hat{m}_i = m_i/m_B$ and

$$\hat{P}_P = 2\sqrt{s}|\mathbf{p}_P|/m_B = \sqrt{(1-s-\hat{m}_P^2)^2 - 4s\hat{m}_P^2}. \quad (17)$$

Since the differential decay rate in Eq. (16) involves the polar angle of the lepton, we can define an angular asymmetry to be

$$\mathcal{A}(q^2) = \frac{\int_{-1}^1 dz \text{sign}(z) d\Gamma_P/(dq^2 dz)}{\int_{-1}^1 dz d\Gamma_P/(dq^2 dz)} \quad (18)$$

with $z = \cos\theta$. Explicitly, the asymmetry for $\bar{B} \rightarrow P\ell\bar{\nu}_\ell$ decay is

$$\mathcal{A}_P(s) = \frac{\Gamma_2^P}{2\Gamma_1^P + 2/3\Gamma_3^P}. \quad (19)$$

Moreover, the integrated angular asymmetry can be defined by

$$\bar{\mathcal{A}}_P = \frac{\int dq^2 \int_{-1}^1 dz \text{sign}(z) d\Gamma_P/(dq^2 dz)}{\int dq^2 \int_{-1}^1 dz d\Gamma_P/(dq^2 dz)}. \quad (20)$$

The angular asymmetry is only associated with the ratio of form factors, which supposedly is insensitive to the hadronic parameters. Plausibly, this physical quantity could be the good candidate to explore the new physics such as charged Higgs [14], right-handed gauge boson [15], etc.

Before presenting the numerical results for the form factors and other related quantities, we will briefly discuss how to extract the input parameters for the η_q in the presence of $\eta - \eta'$

mixing. Following the divergences of the axial vector currents

$$\partial^\mu \bar{q}' \gamma_\mu \gamma_5 q' = \frac{\alpha_s}{4\pi} G \tilde{G} + 2m_{q'} \bar{q}' i \gamma_5 q', \quad (21)$$

where $G = G^{a\mu\nu}$ are the gluonic field-strength and $\tilde{G} = \tilde{G}^{a\mu\nu} \equiv \epsilon^{\mu\nu\alpha\beta} G_{\alpha\beta}^a$, the mass matrix of $\eta_{q,s}$ becomes

$$\begin{aligned} \begin{pmatrix} M_{qq}^2 & M_{qs}^2 \\ M_{sq}^2 & M_{ss}^2 \end{pmatrix} &= \begin{pmatrix} \langle 0 | \partial^\mu J_{\mu 5}^q | \eta_q \rangle / f_q & \langle 0 | \partial^\mu J_{\mu 5}^s | \eta_q \rangle / f_s \\ \langle 0 | \partial^\mu J_{\mu 5}^q | \eta_s \rangle / f_q & \langle 0 | \partial^\mu J_{\mu 5}^s | \eta_s \rangle / f_s \end{pmatrix} \\ &= \begin{pmatrix} m_{qq}^2 + 2a^2 & \sqrt{2}ya^2 \\ \sqrt{2}ya^2 & m_{ss}^2 + y^2a^2 \end{pmatrix} \end{aligned} \quad (22)$$

with $a^2 = \langle 0 | \alpha_s G \tilde{G} | \eta_q \rangle / (4\sqrt{2}\pi f_q)$ and $y = f_q / f_s$. Using the mixing matrix introduced in Eq. (1), one can diagonalize the mass matrix and the eigenvalues are the physical mass of η and η' . Correspondingly, we have the relations [13]

$$\begin{aligned} \sin \phi &= \left[\frac{(m_{\eta'}^2 - m_{ss}^2)(m_\eta^2 - m_{qq}^2)}{(m_{\eta'}^2 - m_\eta^2)(m_{ss}^2 - m_{qq}^2)} \right]^{1/2}, \\ y &= \left[2 \frac{(m_{\eta'}^2 - m_{ss}^2)(m_{ss}^2 - m_\eta^2)}{(m_{\eta'}^2 - m_{qq}^2)(m_\eta^2 - m_{qq}^2)} \right]^{1/2}, \\ a^2 &= \frac{1}{2} \frac{(m_{\eta'}^2 - m_{qq}^2)(m_\eta^2 - m_{qq}^2)}{m_{ss}^2 - m_{qq}^2}, \end{aligned} \quad (23)$$

and $m_{\eta^{(\prime)}}$ is the mass of $\eta^{(\prime)}$. Once the parameters ϕ , y and a are determined by experiments, we can get the information for $m_{qq,ss}$ and $f_{q,s}$. Then, they could be taken as the inputs in our calculations.

After formulating the necessary pieces, we now perform the numerical analysis for the form factors and the related physical quantities introduced earlier. For understanding how well the predictions of LF model are, we first analyze $B \rightarrow \pi$ form factors at $q^2 = 0$. By examining Eq. (11), we see that the main theoretical unknowns are the parameters of distribution amplitudes of mesons, masses of constitute quarks and the decay constants of mesons. As usual, we adopt the gaussian-type wave function for pseudoscalar mesons as

$$\varphi'_P(x_2, p'_\perp) = 4 \left(\frac{\pi}{\beta_P'^2} \right)^{\frac{3}{4}} \sqrt{\frac{dp'_z}{dx_2}} \exp \left(-\frac{p_z'^2 + p_\perp'^2}{2\beta_P'^2} \right), \quad (24)$$

with β'_P characterizing the shape of the wave function. Other relevant values of parameters are taken as (in units of GeV)

$$\begin{aligned} m_B &= 5.28, \quad m_b = (4.8 \pm 0.2), \quad m_\pi = 0.14, \\ m_u &= m_d = (0.26 \pm 0.03), \quad f_B = (0.19 \pm 0.02), \quad f_\pi = 0.131, \\ \beta'_B &= 0.553^{+0.047}_{-0.048}, \quad \beta'_\pi = 0.31, \quad \beta'_{\eta_q} = 0.353^{+0.014}_{-0.013}, \end{aligned} \quad (25)$$

where $m_{u,d}$ are the constituent quark masses, the errors in them are from the combination of linear, harmonic oscillator and power law potential [16] and f_P denotes the decay constant of P-meson. The shape parameters βs are determined by the relevant decay constants whose analytic expressions are given in Ref. [10]. Following the formulae derived in Eq. (11) and using the taken values of parameters, we immediately find

$$f_+^\pi(0) = 0.245^{+0.000+0.011}_{-0.001-0.011}, \quad f_T^\pi(0) = 0.239^{+0.002+0.020}_{-0.003-0.018}, \quad (26)$$

where the two kinds of uncertainties are from (i) β'_B ; (ii) the quark masses m_u and m_b (added in quadrature). To compare with the results of LCSRs given by $f_+^\pi(0)|_{LCSR} = 0.258 \pm 0.031$ and $f_T^\pi(0)|_{LCSR} = 0.253 \pm 0.028$ [17], it is clear that although the central value of LF model is slightly smaller than those of LCSRs, they are still consistent with each other by counting the errors. Since we use the quark-flavor scheme, for estimating the form factors associated with $\eta^{(\prime)}$, the values of involving parameters are chosen to be $f_q = (1.07 \pm 0.02)f_\pi$, $\phi = 39.3^\circ \pm 1.0^\circ$ [12], $m_{qq} = 0.14^{+0.11}_{-0.04}$ GeV [7], $f_{+(T)}^\eta = \cos \phi f_{+(T)}^{\eta_q}$ and $f_{+(T)}^{\eta'} = \sin \phi f_{+(T)}^{\eta_q}$, we have

$$\begin{aligned} f_+^{\eta_q}(0) &= 0.284^{+0.012+0.012}_{-0.012-0.011}, \quad f_T^{\eta_q}(0) = 0.273^{+0.011+0.022}_{-0.011-0.020}, \\ f_+^\eta(0) &= 0.220^{+0.009+0.009}_{-0.009-0.009}, \quad f_T^\eta(0) = 0.211^{+0.009+0.017}_{-0.009-0.015}, \\ f_+^{\eta'}(0) &= 0.180^{+0.008+0.008}_{-0.008-0.007}, \quad f_T^{\eta'}(0) = 0.173^{+0.007+0.014}_{-0.007-0.013}, \end{aligned} \quad (27)$$

where the first and second errors are from (i) β'_B and β'_{η_q} (ii) the quark masses m_u and m_b , respectively. From Eq. (11), one can see that the form factor $f_+^{\eta_q}(q^2)$ does not depend on the mass m_{qq} , while the dependence of m_{qq} in $f_T^{\eta_q}$ resides in the term $M' + M''$ (in this case $m_B + m_{qq}$). The uncertainty of f_T caused by the m_{qq} is less than 2%. Furthermore, since the form factors are associated with mixing angle ϕ , the corresponding uncertainties for $B \rightarrow \eta^{(\prime)}$ and BRs of $\bar{B} \rightarrow \eta^{(\prime)} \ell \bar{\nu}_\ell$ are expected to be 2.1% (1.4%) and 4.2% (2.8%), respectively. Despite

different treatments of quarks' momenta, the results here are well consistent with that in light-cone quark model constructed in the effective field theory [18]: $f_+^{\eta q}(0) = 0.287_{-0.065}^{+0.059}$. Intriguingly, our results are also consistent with $f_+^{\eta}(0)|_{LCSR} = 0.231_{-0.020}^{+0.018}$ and $f_+^{\eta'}(0)|_{LCSR} = 0.189_{-0.016}^{+0.015}$ calculated by LCSRs [19]. In order to understand the behavior of whole q^2 , the form factors for $B \rightarrow P$ are parametrized by [17]

$$F_i(q^2) = \frac{F_i(0)}{1 - aq^2/m_B^2 + b(q^2/m_B^2)^2}, \quad (28)$$

where F_i denotes any form factor among $f_{+,0,T}$. The fitted values of a, b for $B \rightarrow (\pi, \eta, \eta')$ are displayed in Table I, where the uncertainties are similar to the ones given in Eq. (27).

TABLE I: Values of parameters for q^2 dependent $B \rightarrow (\pi, \eta, \eta')$ form factors calculated by LF quark model.

$F(q^2)$	a	b	$F(q^2)$	a	b
$f_+^{B \rightarrow \pi}$	$1.62_{-0.05-0.11}^{+0.05+0.12}$	$0.79_{-0.08-0.15}^{+0.09+0.17}$	$f_+^{B \rightarrow \eta^{(\prime)}}$	$1.55_{-0.05-0.10}^{+0.04+0.11}$	$0.65_{-0.06-0.12}^{+0.08+0.13}$
$f_0^{B \rightarrow \pi}$	$0.75_{-0.03-0.10}^{+0.04+0.11}$	$0.07_{-0.03-0.05}^{+0.02+0.05}$	$f_0^{B \rightarrow \eta^{(\prime)}}$	$0.67_{-0.03-0.09}^{+0.03+0.09}$	$0.03_{-0.02-0.03}^{+0.03+0.03}$
$f_T^{B \rightarrow \pi}$	$1.60_{-0.05-0.11}^{+0.05+0.12}$	$0.75_{-0.08-0.15}^{+0.09+0.17}$	$f_T^{B \rightarrow \eta^{(\prime)}}$	$1.53_{-0.04-0.10}^{+0.04+0.11}$	$0.62_{-0.07-0.11}^{+0.08+0.13}$

In the quark flavor mixing mechanism, the η and η' meson receives additional coupling with two gluons, due to the axial anomaly. Thus to be self-consistent, in the study of the transition form factors, one also needs to include the so-called gluonic form factors which is induced by the transition from the two gluons into the $\eta^{(\prime)}$. In our study, the gluonic form factors have been neglected and there are two reasons for this. In the light-front quark model, the leading order contribution to the form factor is of the order α_s^0 while the gluonic form factor is suppressed by the α_s , where the coupling constant is evaluated at the typical scale $\mu \sim \sqrt{\Lambda_{\text{QCD}} \times m_B}$ (with Λ_{QCD} hadronic scale). The inclusion of the gluonic form factors also requires the next-to-leading order studies for the quark content, which is beyond the scope of the present work. Secondly the factorization analysis of the gluonic form factors such as the perturbative QCD study in Ref. [22] reflects that there is no endpoint singularity in the gluonic form factors and the PQCD study shows that the gluonic form factors are negligibly small. This feature is also confirmed by the recent LCSR results [19]. For terms without endpoint singularity, different approaches usually obtain similar results. Thus our results of the semileptonic $B \rightarrow \eta^{(\prime)} l \bar{\nu}$ will not be sizably affected by the gluonic form factors,

although they are not taken into account in the present analysis.

Besides the form factors could be the source of uncertainties, another uncertain quantity in exclusive $b \rightarrow u\ell\bar{\nu}_\ell$ decays is from the Cabibbo-Kobayashi-Maskawa (CKM) matrix element $V_{ub} \sim \lambda^3$ with λ being Wolfenstein parameter. Results for V_{ub} determined by inclusive and exclusive decaying modes have some inconsistencies [15, 20]. For a self-consistent analysis, we take $B \rightarrow \pi$ form factors calculated by LF model and the data $\mathcal{B}(\bar{B}_d \rightarrow \pi^+\ell'\bar{\nu}_{\ell'}) = (1.36 \pm 0.09) \times 10^{-4}$ with $\ell' = e, \mu$ [20] as the inputs to determine the $|V_{ub}|$. Neglecting the lepton mass, one gets the differential decaying rate for $\bar{B} \rightarrow \pi\ell'\bar{\nu}_{\ell'}$

$$\frac{d\Gamma_\pi}{dq^2} = \frac{G_F^2 |V_{ub}|^2 m_B^3}{3 \cdot 2^6 \pi^3} \sqrt{(1-s+\hat{m}_\pi^2)^2 - 4\hat{m}_\pi^2} \left(f_+^\pi(q^2) \hat{P}_\pi \right)^2, \quad (29)$$

where only the f_+^π form factor involves. Accordingly, the value of V_{ub} is found by

$$|V_{ub}|_{LF} = (3.99 \pm 0.13) \times 10^{-3}. \quad (30)$$

With the obtained result of $|V_{ub}|_{LF}$, the form factors in the Table I, the predicted BRs for $B^- \rightarrow (\eta, \eta')\ell\bar{\nu}_\ell$, together with the experimental results measured by BaBar collaboration [21], are displayed in Table II. The predicted result for the BR of $B^- \rightarrow \eta\ell\bar{\nu}_\ell$ is about two times larger than that of $B^- \rightarrow \eta'\ell\bar{\nu}_\ell$: the form factor of $B^- \rightarrow \eta$ is larger than the form factor of $B^- \rightarrow \eta'$; the phase space in $B \rightarrow \eta'\ell\bar{\nu}_\ell$ is smaller. Branching ratios for decays with a tau lepton are naturally smaller than the relevant channels with a lighter lepton.

TABLE II: BRs of $B^- \rightarrow \eta^{(\prime)}\ell\bar{\nu}_\ell$ (in units of 10^{-4}). The two kinds of errors shown in the table are from (i) $\beta'_B, \beta'_{\eta_q}$; (ii) the quark masses m_u, m_b , respectively.

Mode	$B^- \rightarrow \eta\ell'\bar{\nu}_{\ell'}$	$B^- \rightarrow \eta\tau\bar{\nu}_\tau$	$B^- \rightarrow \eta'\ell'\bar{\nu}_{\ell'}$	$B^- \rightarrow \eta'\tau\bar{\nu}_\tau$
This work	$0.49^{+0.02+0.10}_{-0.04-0.07}$	$0.29^{+0.01+0.07}_{-0.02-0.05}$	$0.24^{+0.01+0.04}_{-0.02-0.03}$	$0.13^{+0.01+0.03}_{-0.01-0.02}$
Exp. [21]	$0.31 \pm 0.06 \pm 0.08$		< 0.47	

According to Eq. (19), moreover, we can study the lepton angular asymmetries. Using the obtained form factors, we present the asymmetry as a function of q^2 in Fig. 2, where the solid, dashed and dash-dotted lines are for $B^- \rightarrow (\pi^0, \eta, \eta')\tau\bar{\nu}_\tau$, respectively. Due to the angular asymmetry being proportional to m_ℓ^2 in the SM, here we only present the effects on τ decaying modes. At very small q^2 region, the three lines are approaching the point 0.75 which can be easily derived from the definition of angular asymmetries. At the small

recoil region, the $\hat{P}_P$ defined in the Eq. (17) is approaching zero and all the lepton angular asymmetries are close to 0. The integrated angular asymmetries defined in Eq. (20) for $B^- \rightarrow (\pi^0, \eta, \eta')\tau\bar{\nu}_\tau$ are predicted by $(0.277^{+0.001+0.005}_{-0.001-0.007}, 0.290^{+0.002+0.003}_{-0.000-0.003}, 0.312^{+0.004+0.005}_{-0.000-0.006})$.




FIG. 2: Angular asymmetries for $B \rightarrow (\pi, \eta, \eta')\ell\nu_\ell$ are depicted by the solid (black), dashed (red) and dash-dotted (blue) lines, respectively.

Finally, we make some remark on the D decays. We find that the obtained information on $\eta^{(\prime)}$ can be directly applied to the semileptonic $D^+ \rightarrow \eta^{(\prime)}\ell\nu_\ell$ decays. Since the associated CKM matrix element $|V_{cd}| = 0.2256 \pm 0.0010$ has small errors, if the decay constant of D -meson is well controlled, the $\eta^{(\prime)}$ production in D decays could be the good environment to test the properties of $\eta^{(\prime)}$. Hence, by similar calculations performed in B decays and taking $f_D = (0.205 \pm 0.020)$ GeV, $\beta_D = (0.462^{+0.048}_{-0.047})$ GeV and $m_c = (1.4 \pm 0.1)$ GeV, the form factors for $f_{+,0,T}^{D \rightarrow \eta^{(\prime)}}(q^2)$ are obtained by

$$\begin{aligned} f_+^{D \rightarrow \eta_q}(q^2) &= \frac{0.688}{1 - 1.03\hat{s} + 0.29\hat{s}^2}, \\ f_0^{D \rightarrow \eta_q}(q^2) &= \frac{0.705}{1 - 0.39\hat{s} + 0.01\hat{s}^2}, \\ f_T^{D \rightarrow \eta_q}(q^2) &= \frac{0.616}{1 - 1.08\hat{s} + 0.25\hat{s}^2}, \end{aligned} \quad (31)$$

where we have used the parametrization defined in Eq. (28), $\hat{s} = q^2/m_D^2$ and only the central values are shown. The small differences between $f_+(0)$ and $f_0(0)$ arise from the fitting procedure. Replacing the parameters of B -meson appearing in Eq. (29) by those of D -meson, the BRs for $D^- \rightarrow (\eta, \eta')\ell\bar{\nu}_\ell$ are predicted by

$$\begin{aligned} \mathcal{B}(D^- \rightarrow \eta\ell\bar{\nu}_\ell) &= (1.11^{+0.05+0.09}_{-0.06-0.09}) \times 10^{-3}, \\ \mathcal{B}(D^- \rightarrow \eta'\ell\bar{\nu}_\ell) &= (1.79^{+0.07+0.12}_{-0.08-0.12}) \times 10^{-4}, \end{aligned} \quad (32)$$

respectively. It is found that $\mathcal{B}(D^- \rightarrow \eta' \ell \bar{\nu}_\ell)$ is almost one order of magnitude smaller than $\mathcal{B}(D^- \rightarrow \eta \ell \bar{\nu}_\ell)$. The reason for the resulted smallness is just phase space suppression. Our predictions are well consistent with the recent measurements by the CLEO collaboration [23]:

$$\begin{aligned}\mathcal{B}(D^- \rightarrow \eta \ell \bar{\nu}_\ell) &= (1.33 \pm 0.20 \pm 0.06) \times 10^{-3}, \\ \mathcal{B}(D^- \rightarrow \eta' \ell \bar{\nu}_\ell) &< 3.5 \times 10^{-4}.\end{aligned}\tag{33}$$

This consistence is very encouraging. The $D^- \rightarrow \eta' l \bar{\nu}$ may be detected in the near future. Our results are also consistent with the results given in Ref. [24].

In summary, we have calculated the $B \rightarrow (\pi, \eta, \eta')$ transition form factors in LF approach. We find that at maximum recoil the values of form factors are $f_+^{(\pi, \eta, \eta')}(0) = (0.245_{-0.001}^{+0.000} \pm 0.011, 0.220 \pm 0.009 \pm 0.009, 0.180 \pm 0.008_{-0.007}^{+0.008})$ and $f_T^{(\pi, \eta, \eta')}(0) = (0.239_{-0.003-0.018}^{+0.002+0.020}, 0.211 \pm 0.009_{-0.015}^{+0.017}, 0.173 \pm 0.007_{-0.013}^{+0.014})$, respectively. Our calculated values are consistent with the results done by LCSRs. With the obtained form factor $f_+^\pi(q^2)$ and observed BR for $\bar{B}_d \rightarrow \pi^+ \ell \bar{\nu}_\ell$, the V_{ub} is extracted to be $|V_{ub}|_{LF} = (3.99 \pm 0.13) \times 10^{-3}$. Accordingly, we predict the BRs for $\bar{B} \rightarrow (\eta, \eta') \ell \bar{\nu}_\ell$ decays with $\ell = e, \mu$ as $(0.49_{-0.04-0.07}^{+0.02+0.10}, 0.24_{-0.02-0.03}^{+0.01+0.04}) \times 10^{-4}$, while the BRs for $D^- \rightarrow (\eta, \eta') \ell \bar{\nu}_\ell$ are given by $(11.1_{-0.6-0.9}^{+0.5+0.9}, 1.79_{-0.08-0.12}^{+0.07+0.12}) \times 10^{-4}$. In addition, we also show that the lepton angular asymmetries for $\bar{B} \rightarrow (\pi, \eta, \eta') \tau \bar{\nu}_\tau$ are $(0.277_{-0.001-0.007}^{+0.001+0.005}, 0.290_{-0.000-0.003}^{+0.002+0.003}, 0.312_{-0.000-0.006}^{+0.004+0.005})$.

Acknowledgments

This work is supported by the National Science Council of R.O.C. under Grant No: NSC-97-2112-M-006-001-MY3, and National Natural Science Foundation of China under the Grant No. 10847161.

-
- [1] CLEO Collaboration, T.E. Browder *et al.*, Phys. Rev. Lett. **81**, 1786 (1998).
 - [2] E. Barberio *et al.*, arXiv:0808.1297, online update at <http://www.slac.stanford.edu/xorg/hfag>.
 - [3] I. Halperin and A. Zhitnitsky, Phys. Rev. D **56**, 7247 (1997); A.A. Petrov, Phys. Rev. D **58**, 054004 (1999); T. Feldmann and P. Kroll, Eur. Phys. J. C **5**, 327 (1998); T. Feldmann, P. Kroll and B. Stech, Phys. Lett. B **449**, 339 (1999).

- [4] D. Atwood and A. Soni, Phys. Lett. B**405**, 150 (1997); M.R. Ahmady, *et al.*, Phys. Rev. D**58**, 014015 (1998); M.R. Ahmady and E. Kou, Phys. Rev. D**59**, 054014 (1999).
- [5] M. Z. Yang and Y. D. Yang, Nucl. Phys. B**609**, 469 (2001).
- [6] M. Beneke and M. Neubert, Nucl. Phys. B**651**, 225 (2003).
- [7] C. H. Chen and C. Q. Geng, Phys. Lett. B**645**, 197 (2007); A. Akeroyd, C. H. Chen and C. Q. Geng, Phys. Rev. D**75**, 054003 (2007).
- [8] Y. Y. Keum, H. n. Li and A. I. Sanda, Phys. Lett. B **504**, 6 (2001); Phys. Rev. D **63**, 054008 (2001); C. D. Lu, K. Ukai and M. Z. Yang, Phys. Rev. D **63**, 074009 (2001).
- [9] W. Jaus, Phys. Rev. D **60**, 054026 (1999).
- [10] H. Y. Cheng, C. K. Chua and C. W. Hwang, Phys. Rev. D**69**, 074025 (2004).
- [11] J. Schechter, A. Subbaraman and H. Weigel, Phys. Rev. D **48**, 339 (1993).
- [12] T. Feldmann, P. Kroll, B. Stech, Phys. Rev. D**58**, 114006(1998).
- [13] T. Feldmann and P. Kroll, Phys. Scripta T**99**, 13 (2002).
- [14] C. H. Chen and C. .Q. Geng, Phys. Rev. D**71**, 077501 (2005); JHEP**0610**, 053 (2006).
- [15] C. H. Chen and S. H. Nam, Phys. Lett. B**666**, 462 (2008).
- [16] H. M. Choi, arXiv:hep-ph/9911271.
- [17] P. Ball and R. Zwicky, Phys. Rev. D**71**, 014015 (2005).
- [18] C. D. Lu, W. Wang and Z. T. Wei, Phys. Rev. D **76**, 014013 (2007).
- [19] P. Ball and G. W. Jones, JHEP**0708**, 025 (2007).
- [20] C. Amsler *et al.* (Particle Data Group), Phys. Lett. B**667**, 1 (2008).
- [21] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **101**, 081801; Phys. Rev. D**79**, 052011 (2009).
- [22] Y. Y. Charng, T. Kurimoto and H. n. Li, Phys. Rev. D **74**, 074024 (2006) [Phys. Rev. D **78**, 059901 (2008)].
- [23] R. E. Mitchell *et al.* [CLEO Collaboration], Phys. Rev. Lett. **102**, 081801 (2009).
- [24] Z. T. Wei, H. W. Ke and X. F. Yang, Phys. Rev. D **80**, 015022 (2009) [arXiv:0905.3069 [hep-ph]].